


Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Journal of Hydrology

journal homepage: [www.elsevier.com/locate/jhydrol](http://www.elsevier.com/locate/jhydrol)

## Discussion

Comment on “Interference of river level changes on riparian zone evapotranspiration estimates from diurnal groundwater level fluctuations” by J. Zhu, M. Young, J. Healy, R. Jasoni, J. Osterberg [J. Hydrol. 403(3–4) (2011) 381–389]

Jozsef Szilagyi <sup>a,b,\*</sup>, Zoltan Gribovszki <sup>c</sup>, Peter Kalicz <sup>c</sup>

<sup>a</sup> Department of Hydraulic and Water Resources Engineering, Budapest University of Technology and Economics, Budapest H-1111, Hungary

<sup>b</sup> School of Natural Resources, University of Nebraska-Lincoln, Lincoln 68583, USA

<sup>c</sup> Institute of Geomatics and Civil Engineering, University of Western Hungary, Sopron H-9400, Hungary




Fig. 1. The experimental catchment and the location of the groundwater wells [after Gribovszki et al. (2008)].

Zhu et al. draw the attention to the importance of considering stream water level fluctuations as a complicating factor in estimating diurnal evapotranspiration rates by riparian/phreatophyte vegetation from groundwater level fluctuations. To justify their analysis they cite the studies of Gribovszki et al. (2008) and Szilagyi et al. (2008) among other works. While Zhu et al.'s analysis


of the interaction of streamwater and groundwater level fluctuations is definitely justified for streams that are influenced by streamwater/groundwater pumping and/or flow releases (the Colorado River below Hoover Dam is a good example), we would like to point out that in smaller streams unaffected by human interference, such interactions may largely be negligible as diurnal water level changes in the stream are significantly damped in comparison with the corresponding groundwater level changes, as can be demonstrated in the watershed discussed by Gribovszki et al. (2008).

The Hidegvíz Valley catchment (Fig. 1) is a small forested watershed (6 km<sup>2</sup>) within the hilly sub-alpine terrain of western Hun-

DOI of original article: [10.1016/j.jhydrol.2011.08.005](https://doi.org/10.1016/j.jhydrol.2011.08.005)

\* Corresponding author at: Department of Hydraulic and Water Resources Engineering, Budapest University of Technology and Economics, Budapest, H-1111, Hungary.

E-mail address: [jszilagyi1@unl.edu](mailto:jszilagyi1@unl.edu) (J. Szilagyi).



**Fig. 2.** Water level changes in the stream (smooth line) and in the groundwater well 9.4 m from the stream (well #2 in Fig. 1). The largest fluctuations are caused by rain events.

gary near the border of Austria. For more detailed characteristics of the site see Gribovszki et al. (2008). The watershed may be a prototype of other small forested catchments of similar physiography under a continental climate. As Fig. 2 illustrates, diurnal streamwater level fluctuations are vastly negligible to the corresponding changes in groundwater level, obtained in a well 9.4 m from the stream. For clarity of presentation, groundwater level fluctuations only in this well are shown, the fluctuations are similar in magnitude in the other well locations of Fig. 1, and all are at least an order of magnitude larger than the corresponding fluctuations in stream water level. This is a welcome news to the application of observed diurnal groundwater level fluctuations for the estimation of riparian/phreatophyte evapotranspiration rates in catchments with no or minor anthropogenic influence, since the resulting estimates are not adversely affected by the distance to the stream, as on the site, discussed by Zhu et al. Without taking

away from the importance of their study, we felt it necessary to point out that the problem they discuss is not general, although probably widespread due to the degree humans interfere with the hydrologic cycle. Yet, there remain as many small watersheds not yet influenced by such activities where similar concerns do not enter into the analysis of diurnal groundwater level fluctuations for the estimation of riparian/phreatophyte vegetation water use, as is the case for the study of Gribovszki et al. (2008).

## References

- Gribovszki, Z., Kalicz, P., Szilagyi, J., Kucsara, M., 2008. Riparian zone evapotranspiration estimation from diurnal groundwater level fluctuations. *J. Hydrol.* 349 (1–2), 6–17.
- Szilagyi, J., Gribovszki, Z., Kalicz, P., Kucsara, M., 2008. On diurnal riparian zone groundwater level and streamflow fluctuations. *J. Hydrol.* 349 (1–2), 1–5.